

# FOR THE ACUTE TREATMENT OF MIGRAINE AND PREVENTIVE TREATMENT OF EPISODIC MIGRAINE IN ADULTS<sup>1</sup>

**Nurtec<sup>®</sup> ODT**  
(rimegepant)  
orally disintegrating tablets 75 mg

## Nurtec<sup>®</sup> ODT (rimegepant) is the only oral CGRP receptor antagonist approved for acute and preventive treatment and covered on Express Scripts' national commercial formularies<sup>2-4,†</sup>

### Formulary coverage for oral CGRP receptor antagonists

DRUG NAME	COVERAGE DETAILS <sup>1,3-5</sup>	
	Acute treatment	Preventive treatment*
<b>Nurtec ODT 75 mg</b>	<ul style="list-style-type: none"><li>• Prior trial of 1 triptan required</li><li>• Covered</li></ul>	<ul style="list-style-type: none"><li>• Prior trial of 2 agents for migraine prophylaxis<sup>‡</sup></li><li>• Covered</li></ul>
<b>Ubrelvy<sup>®</sup> (ubrogepant) 50 mg, 100 mg</b>	<ul style="list-style-type: none"><li>• Prior trial of 1 triptan required</li><li>• Covered</li></ul>	Not indicated

CGRP, calcitonin gene-related peptide; ODT, orally disintegrating tablet.

\*Preventive treatment of episodic migraine only. †Managed Markets Insights & Technology LLC as of 07/21. ‡Prior prophylactic therapies must be from different classes. Examples include angiotensin receptor blockers, angiotensin converting enzyme inhibitors, anticonvulsants, beta-blockers, calcium channel blockers, tricyclic antidepressants, or other depressants.

### INDICATION

Nurtec ODT is indicated in adults for the:

- acute treatment of migraine with or without aura
- preventive treatment of episodic migraine

### IMPORTANT SAFETY INFORMATION

**Contraindications:** Hypersensitivity to Nurtec ODT or any of its components.

**Warnings and Precautions:** If a serious hypersensitivity reaction occurs, discontinue Nurtec ODT and initiate appropriate therapy. Serious hypersensitivity reactions have included dyspnea and rash, and can occur days after administration.

**Adverse Reactions:** The most common adverse reactions were nausea (2.7% in patients who received Nurtec ODT compared to 0.8% in patients who received placebo) and abdominal pain/dyspepsia (2.4% in patients who received Nurtec ODT compared to 0.8% in patients who received placebo). Hypersensitivity, including dyspnea and rash, occurred in less than 1% of patients treated with Nurtec ODT.

**Drug Interactions:** Avoid concomitant administration of Nurtec ODT with strong inhibitors of CYP3A4, strong or moderate inducers of CYP3A or inhibitors of P-gp or BCRP. Avoid another dose of Nurtec ODT within 48 hours when it is administered with moderate inhibitors of CYP3A4.

**Use in Specific Populations:** *Pregnant/breast feeding:* It is not known if Nurtec ODT can harm an unborn baby or if it passes into breast milk. *Hepatic impairment:* Avoid use of Nurtec ODT in persons with severe hepatic impairment. *Renal impairment:* Avoid use in patients with end-stage renal disease.

**Please see accompanying full Prescribing Information.**

**References:** **1.** Nurtec ODT. Package Insert. Biohaven Pharmaceuticals Inc. **2.** Croop L, Lipton RB, Kudrow D, et al. Oral rimegepant for preventive treatment of migraine: a phase 2/3, randomised, double-blind, placebo-controlled trial. *Lancet*. Published online December 15, 2020. doi:10.1016/S0140-6736(20)32544-7. **3.** Express Scripts website. Migraine-Nurtec ODT prior authorization policy. <https://corporate-site-labs-prod.s3.us-east-2.amazonaws.com/2021-08/Migraine%20-%20Nurtec%20ODT%20PA%20Policy.pdf>. Updated July 21, 2021. Accessed September 2, 2021. **4.** Express Scripts website. Migraine-Ubrelvy prior authorization policy. <https://corporate-site-labs-prod.s3.us-east-2.amazonaws.com/2021-05/Migraine%20-%20Ubrelvy%20PA%20Policy.pdf>. Updated February 3, 2021. Accessed September 2, 2021. **5.** Ubrelvy. Package insert. Allergan, Inc.

© 2021 Biohaven Pharmaceuticals, Inc. All rights reserved. NURTEC and NURTEC ODT are registered trademarks of Biohaven Pharmaceutical Ireland DAC. Biohaven is a trademark of Biohaven Pharmaceutical Holding Company Ltd.

US-RIMODT-2100786 09/02/2021

