

# Treating Migraine via Telemedicine



## Tips for Prescribing Nurtec ODT Virtually

Set Nurtec ODT (NDC: 72618-3000-2) as a favorite option in your EHR for quick access in two easy steps<sup>1</sup>:

**1 Draft a properly formatted prescription in your EHR.**

**Refer to these helpful prescribing tips:**

- NURTEC ODT 75 mg
- Once daily PRN for migraine
- Dispense #8

**Additional prescribing considerations<sup>1</sup>:**

**Packaging:** Nurtec ODT comes in 8-tablet packs

**Once daily PRN:** Take one 75 mg orally disintegrating tablet (ODT), as needed, for migraine

**24 hour max:** one 75 mg ODT

**30 day max:** The safety of treating more than 15 migraines in a 30-day period has not been established

**2 Save this prescription as a favorite option.**

Use the following chart to find your EHR’s specific term for favorite:

EHR Platform	Term for Favorites
Allscripts Professional EHR	Short List
Cerner	Favorite
Epic	Preference
Practice Fusion	Frequent Medications

This two-step process should help make e-prescribing Nurtec ODT simple. Consider asking your patients if they would like you to send their prescription to an online pharmacy.

### SELECT IMPORTANT SAFETY INFORMATION

**Contraindications:** Hypersensitivity to Nurtec ODT or any of its components.

Please see additional Important Safety Information on the next page and click here for full [Prescribing Information](#).

## SAVINGS AND SUPPORT FOR YOU AND YOUR PATIENTS



**Prior Authorization** assistance available through **covermymeds**<sup>®</sup>  
[www.covermymeds.com](http://www.covermymeds.com)  
 1-866-452-5017



**Nurtec ODT Patient Support Program\*** committed to helping your eligible patients afford their medication.  
[www.nurtec-hcp.com/savings-support](http://www.nurtec-hcp.com/savings-support)  
 \*For full Terms & Conditions, see site above

## IMPORTANT SAFETY INFORMATION

**Contraindications:** Hypersensitivity to Nurtec ODT or any of its components.

**Warnings and Precautions:** If a serious hypersensitivity reaction occurs, discontinue Nurtec ODT and initiate appropriate therapy. Serious hypersensitivity reactions have included dyspnea and rash, and can occur days after administration.

**Adverse Reactions:** The most common adverse reaction was nausea (2% in patients who received Nurtec ODT compared to 0.4% in patients who received placebo). Hypersensitivity, including dyspnea and rash, occurred in less than 1% of patients treated with Nurtec ODT.

**Drug Interactions:** Avoid concomitant administration of Nurtec ODT with strong inhibitors of CYP3A4, strong or moderate inducers of CYP3A or inhibitors of P-gp or BCRP. Avoid another dose of Nurtec ODT within 48 hours when it is administered with moderate inhibitors of CYP3A4.

**Use in Specific Populations:** *Pregnant/breast feeding:* It is not known if Nurtec ODT can harm an unborn baby or if it passes into breast milk. *Hepatic impairment:* Avoid use of Nurtec ODT in persons with severe hepatic impairment. *Renal impairment:* Avoid use in patients with end-stage renal disease.

## INDICATION

Nurtec ODT is indicated for the acute treatment of migraine with or without aura in adults.

### Limitations of Use

Nurtec ODT is not indicated for the preventive treatment of migraine.

**Please click here for full [Prescribing Information](#).**

Price disclosure information for prescribers is available [here](#).

## REFERENCE:

1. Nurtec ODT. Package insert. Biohaven Pharmaceuticals Inc.

